

CARE VETS

carevets.co.uk

Address:
84 Eastham Village Road
Eastham CH62 0AW

Phone:
(0151) 345 1121

Email:
contactus@carevets.co.uk

DISCHARGE ADVICE

MEDICATIONS

Medications normally start the day after surgery. Any change to this will be discussed on discharge and will be on the medication label. Your pet will likely be sore or tired for the next few days.

ANESTHESIA

Your pet was placed under general anaesthesia as part of their procedure. Anaesthetic drugs can take a number of hours to wear off and may cause some patients, especially older patients, to appear drowsy for 24–48 hours. Please keep your pet warm and dry during this time as the drugs may decrease your pet's body temperature. You may notice a slight cough for up to 48 hours due to irritation of the trachea (windpipe) from placement of the tube during anaesthesia.

Anaesthesia can make your pet vomit if they eats or drinks a large volume after surgery. Please feed half of your pet's diet tonight, and resume normal feeding tomorrow morning. It is normal to notice a decrease in activity or appetite for 1 or 2 days as well as lack of a bowel movement for 24–36 hours after surgery. If you have any questions please to contact the hospital.

INTRAVENOUS CATHETER

Your pet may have had an IV (intravenous) IV catheter placed on one of their legs - if so there will be a bandage on the leg. A shaved area on one of and some mild bruising is normal. The bandage can be removed about half an hour after returning home.

RESTRICTED ACTIVITY

Please restrict your pet's activity for the next 10–14 days. So no walking off-lead, running, jumping. If they sleep upstairs or sit on furniture please carry or lift them, and if there are other pets at home, do not allow play.

E-COLLAR

When discharged with an e-collar please try keep the collar on at all times and keep it clean and dry. Your pet can eat and drink while wearing it.

SUTURE CARE

Sutures are normally under the skin - just because your pet cant see them it doesn't mean that a determined pet cant remove them! Please don't allow your pet to lick the incision until it is healed.

Keep a close eye on the incision - look for swelling, redness and discharge, as these may be signs of infection or a reaction to the suture material. Please call the practice if you have any concerns.



Our arms around your BestFriend.